

Mathematics - Course 221

DIFFERENTIATING EXPONENTIAL FUNCTIONS

I Derivative of $e^{g(x)}$

Recall (lesson 221.20-2) that the derivative of the function $f(x)$ is the 'instantaneous' rate of change of $f(x)$ with respect to x .




Figure 1

In Figure 1, the 'instantaneous' R/C $f(x) = e^{g(x)}$ wrt x at $x = x_1$ is equivalent to

- (1) $\lim_{Q \rightarrow P_1} (\text{slope of secant } P_1Q)$
- (2) slope of tangent P_1T
- (3) $f'(x_1)$, the derivative of $e^{g(x)}$ evaluated at $x = x_1$.

Recall (lesson 221.20-2) the basic defining equation of the derivative of $f(x)$:

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

Applying this equation to $f(x) = e^{g(x)}$ yields

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{e^{g(x + \Delta x)} - e^{g(x)}}{\Delta x}$$

It can be shown (but is beyond the scope of this course to do so) that the above limit reduces to

$$e^{g(x)} g'(x)$$

Hence the formula for the *derivative of an exponential function* is

$$\frac{d}{dx} e^{g(x)} = e^{g(x)} g'(x)$$

Example 1

$$\begin{aligned} \frac{d}{dx} e^x &= e^x \frac{d}{dx} x \\ &= e^x \end{aligned}$$

Note that e^x equals its own derivative!

Example 2

$$\begin{aligned} \frac{d}{dx} 6e^{x^2} &= 6 \frac{d}{dx} e^{x^2} \\ &= 6e^{x^2} \frac{d}{dx} x^2 \\ &= 6e^{x^2} (2x) \\ &= 12xe^{x^2} \end{aligned}$$

Example 3

$$\begin{aligned}
 \frac{d}{dx} e^{2\sqrt{x}} &= e^{2\sqrt{x}} \frac{d}{dx} 2\sqrt{x} \\
 &= e^{2\sqrt{x}} \left(2 \frac{d}{dx} x^{1/2} \right) \\
 &= e^{2\sqrt{x}} (2) \left(\frac{1}{2} x^{-1/2} \right) \\
 &= \frac{e^{2\sqrt{x}}}{\sqrt{x}}
 \end{aligned}$$

Example 4

$$\begin{aligned}
 \frac{d}{dx} (15x^3 - e^{-ax^2}) &= \frac{d}{dx} 15x^3 - \frac{d}{dx} e^{-ax^2} \\
 &= 15 \frac{d}{dx} x^3 - e^{-ax^2} \frac{d}{dx} (-ax^2) \\
 &= 15(3x^2) - e^{-ax^2} \left(-a \frac{d}{dx} x^2 \right) \\
 &= 45x^2 + 2axe^{-ax^2}
 \end{aligned}$$

Example 5

Given the displacement function

$$s(t) = 5t^2 + 100e^{-0.4t},$$

- (a) find the velocity function $v(t)$
- (b) find the acceleration function $a(t)$
- (c) sketch the graphs of $s(t)$, $v(t)$ and $a(t)$ over the interval $0 \leq t \leq 10$

Solution

$$(a) \quad v(t) = s'(t)$$

$$= \frac{d}{dt} (5t^2 + 100e^{-0.4t})$$

$$= 10t + 100e^{-0.4t} \frac{d}{dt} (-0.4t)$$

$$= \underline{\underline{10t - 40e^{-0.4t}}}$$

$$(b) \quad a(t) = \frac{dv}{dt}$$

$$= 10 \frac{d}{dt} t - 40 \frac{d}{dt} e^{-0.4t}$$

$$= 10 - 40e^{-0.4t} \frac{d}{dt} (-0.4t)$$

$$= \underline{\underline{10 + 16e^{-0.4t}}}$$

t	0	1	2	3	4	6	8	10
s	100	72	65	75	100	189	324	502
v	-40	-17	2	18	32	56	78	99
a	26	20.7	17.2	14.8	13.2	11.5	10.7	10.3

The following are sample calculations of those used to produce the above table of values:

$$\begin{aligned} s(10) &= 5(10)^2 + 100e^{-0.4(10)} \\ &= 500 + 100e^{-4} \\ &= 500 + 100(0.018) \\ &= \underline{\underline{501.8}} \end{aligned}$$

$$v(10) = 10(10) - 40e^{-0.4}(10)$$

$$= 100 - 40 (0.018)$$

$$= \underline{\underline{99.3}}$$

$$a(10) = 10 + 16e^{-0.4}(10)$$

$$= 10 + 16 (0.018)$$

$$= \underline{\underline{10.3}}$$

It was stated in lesson 221.20-3 that velocity is the slope of the s-t curve, and that acceleration is the slope of the v-t curve. Are these statements consistent with the curves of Figure 2?

Note that the slope of the s-t curve is negative at $t = 0$, rises to zero at the curve minimum ($t = 1.9$), and then increases positively to $t = 10$. Note that this is precisely the behaviour of the v-t curve.

Note that the v-t curve rises most sharply at $t = 0$, and gradually settles to a slower, almost linear rate of rise. Accordingly one would expect a positive acceleration in the entire interval $0 < t < 10$, and one that would fall from its initial value towards a constant value. This is precisely the behaviour of the a-t curve.




Figure 2

II Application to Nuclear Decay

The number of radioactive atoms remaining in a radioactive source decays exponentially with time, according to the relation.

$$N(t) = N_0 e^{-\lambda t}$$

where $N(t)$ = number of radioactive atoms remaining after t seconds,

N_0 = number of radioactive atoms at time $t = 0$, and

λ is the decay constant of the radionuclide in s^{-1}

To find the R/C N wrt t , ie, the number of nuclei decaying per second, differentiate the above relation wrt time:

$$\begin{aligned} \frac{dN}{dt} &= \frac{d}{dt} N_0 e^{-\lambda t} \\ &= N_0 \frac{d}{dt} e^{-\lambda t} \quad (N_0 \text{ a constant}) \\ &= N_0 e^{-\lambda t} \frac{d}{dt} (-\lambda t) \\ &= N_0 e^{-\lambda t} (-\lambda \frac{d}{dt} t) \\ &= -\lambda N_0 e^{-\lambda t} \end{aligned}$$

$$\therefore \boxed{\frac{dN}{dt} = -\lambda N}$$

Note that $\frac{dN}{dt}$ stands for the rate of increase in N . Hence $\frac{dN}{dt}$ is negative (see minus sign on RHS), since N is actually decreasing.

The number of nuclei decaying per unit time is called the activity of a source.

Example 6

How many radioactive nuclei are required to make a 5 mCi source of a nuclide whose decay constant equals $7.3 \times 10^{-5} \text{ s}^{-1}$? (1 curie = 3.7×10^{10} dps)

Solution

$$\frac{dN}{dt} = -5 \text{ mCi}$$

$$\Rightarrow -\lambda N = -5 \times 10^{-3} \times 3.7 \times 10^{10}$$

$$\therefore N = \frac{5 \times 10^{-3} \times 3.7 \times 10^{10}}{7.3 \times 10^{-5}}$$

$$= \underline{\underline{2.5 \times 10^{12}}}$$

ie, there are 2.5×10^{12} atoms in a 5 mCi Source.

* * *

If source activity is designated "A",

$$\text{then } A(t) = - \frac{dN}{dt} \quad (\text{rate of decrease in } N)$$

$$= \lambda N$$

$$= \lambda N_0 e^{-\lambda t} \quad (\because N = N_0 e^{-\lambda t})$$

$$\text{then } A(0) = \lambda N_0 e^0$$

$$= \lambda N_0$$

Let $A_0 = A(0)$

Then $A_0 = \lambda N_0$

and $A(t) = A_0 e^{-\lambda t}$

ie, the activity $A(t)$ obeys the same exponential relationship as $N(t)$.

Example 7

Find the time required for the activity of a source of decay constant $3.5 \times 10^{-4} \text{ s}^{-1}$ to decay by a factor of 1000.

Solution

Let required time be t_1 .

Then $A(t_1) = A_0 e^{-\lambda t_1}$

ie, $\frac{A(t_1)}{A_0} = e^{-\lambda t_1}$

$\therefore e^{-\lambda t_1} = 0.001 \quad (\because \frac{A(t_1)}{A_0} = \frac{1}{1000})$

Taking natural log of both sides,

$$\ln e^{-\lambda t_1} = \ln 0.001$$

$\therefore -\lambda t_1 = \ln 10^{-3} \quad (\text{cf lesson 321.10-4})$

$\therefore t_1 = \frac{\ln 10^{-3}}{-\lambda}$

$$= \frac{-6.91}{-3.5 \times 10^{-4}}$$

$$= \underline{\underline{2.0 \times 10^4 \text{ seconds or 5.5 hours}}}$$

Example 8

Prove: $t_{1/2} = \frac{0.693}{\lambda}$, where $t_{1/2}$ is the half-life of a radionuclide, ie, the time required for source activity to decay to one-half its original activity.

Solution

$$A(t_{1/2}) = A_0 e^{-\lambda t_{1/2}}$$

$$\therefore \frac{A(t_{1/2})}{A_0} = e^{-\lambda t_{1/2}}$$

$$\therefore e^{-\lambda t_{1/2}} = 0.5 \quad (\because \frac{A(t_{1/2})}{A_0} = 0.5)$$

$$\therefore \ln e^{-\lambda t_{1/2}} = \ln 0.5$$

$$\therefore -\lambda t_{1/2} = -0.693$$

$$\therefore t_{1/2} = \frac{0.693}{\lambda}$$

III Application to Reactor Power Growth

Reactor power grows exponentially in time, approximately according to the relation,

$$P(t) = P_0 e^{\frac{\Delta k}{L} t}$$

where $P(t)$ is reactor power at time t ,

P_0 is reactor power at $t = 0$,

Δk is the reactivity in units of "k",

L is the mean neutron lifetime in the reactor.

For example, if $P_0 = 100$ W, and $\frac{\Delta k}{L} = 0.05$, the graph of $P(t)$ vs t is shown in Figure 3.




Figure 3

DEFINITION: The *reactor period* "T" is the time required for the power to increase by a factor of e.

Proof that Reactor Period = $\frac{L}{\Delta k}$

$$P(T) = eP_0 \quad \text{by definition of } T$$

$$\text{ie,} \quad eP_0 = P_0 e^{\frac{\Delta k}{L} T}$$

$$\therefore \ln e = \ln e^{\frac{\Delta k}{L} T}$$

$$\text{ie,} \quad 1 = \frac{\Delta k}{L} T$$

$$\therefore \boxed{T = \frac{L}{\Delta k}}$$

∴ an alternative form of the power growth equation is

$$P(t) = P_0 e^{t/T}$$

from which it is obvious that each time t increases by T , P increases by a factor of e , consistent with previous definition of T .

Not only does the power $P(t)$ grow exponentially with time, but so also does the rate of growth, $P^1(t)$, as shown below:

$$\begin{aligned} \frac{dP}{dt} &= \frac{d}{dt} P_0 e^{\frac{\Delta k}{L} t} \\ &= P_0 \frac{d}{dt} e^{\frac{\Delta k}{L} t} \\ &= P_0 e^{\frac{\Delta k}{L} t} \frac{d}{dt} \frac{\Delta k}{L} t \\ &= P_0 e^{\frac{\Delta k}{L} t} \frac{\Delta k}{L} \frac{d}{dt} t \\ &= \frac{\Delta k}{L} \underbrace{P_0 e^{\frac{\Delta k}{L} t}}_{P(t)} \end{aligned}$$

$$\therefore \frac{dP}{dt} = \frac{\Delta k}{L} P(t) = \frac{1}{T} P(t)$$

Note that power growth rate $P^1(t)$ is directly proportional to product of reactivity Δk and power $P(t)$. Therefore, given sufficiently high values of Δk and P , P^1 may be so high that rated power is exceeded before the regulation system can arrest power growth.

Thus, for reactor protection, a signal is required to detect dangerously high reactivity values at low power. Such a signal is one whose output varies as the rate of change of the logarithm of reactor power. This signal is known as "rate log power":

$$\begin{aligned}
 \frac{d}{dt} (\ln P(t)) &= \frac{d}{dt} \ln P_0 e^{\frac{\Delta k}{L} t} \\
 &= \frac{d}{dt} \left(\ln P_0 + \ln e^{\frac{\Delta k}{L} t} \right) \\
 &= \frac{d}{dt} \ln P_0 + \frac{d}{dt} \frac{\Delta k}{L} t \\
 &= 0 + \frac{\Delta k}{L} \frac{d}{dt} t \\
 &= \frac{\Delta k}{L}
 \end{aligned}$$

∴ rate log power, $\frac{d}{dt} (\ln P(t)) = \frac{\Delta k}{L} = \frac{1}{T}$

Note that rate log power is proportional to reactivity Δk , independent of reactor power. Hence the reactor can be tripped by this signal at low power, eg, 0.001% full power, long before the *linear rate power*, $P^1(t)$ gets out of hand.

ASSIGNMENT

1. Differentiate:

(a) e^{x^2-4}

(b) $-e^{-x}$

(c) $-e^{-x-1}$

(d) $2e^{-1/\sqrt{x}}$

(e) $5e^{\sqrt{t}} (t^2-1)$

(f) $\frac{1}{3} e^{-1/x^3}$

2. Find (i) $v(t)$ (ii) $a(t)$ (iii) $v(2)$ if

(a) $s(t) = e^t - t^3$

(b) $s(t) = e^{-t} + 2t$

3. Plot $s - t$, $v - t$, $a - t$ curves for the displacement function of 2(a) above over the time interval $0 \leq t \leq 3$. Do the slopes of the $s - t$ and $v - t$ curves appear to verify the definitions, $v(t) = s^1(t)$ and $a(t) = v^1(t)$, respectively?

4. If 2.0×10^{19} radioactive nuclei constitute a 5.0 mCi source, what is the decay constant of the radionuclide? (1 curie = 3.7×10^{10} dps)

5. (a) What is the activity of a source consisting of 7.0×10^{18} radioactive nuclei, and having decay constant $2.4 \times 10^{-4} \text{ s}^{-1}$?

(b) How many radioactive nuclei remain after (i) 20 minutes?
(ii) 6 half-lives?

(c) Calculate the source activity after (i) 20 minutes
(ii) 6 half-lives.


(d) Calculate the half-life of the source.


(e) How long does the source take to decay to 10 mCi?

6. If $N(t) = N_0 e^{-\lambda t}$ and $A = -\frac{dN}{dt}$, prove that (a) $A = \lambda N$

(b) $A(t) = A_0 e^{-\lambda t}$

7. Prove that $t_{1/2} = \frac{\ln 2}{\lambda}$.
8. If $P(t) = P_0 e^{t/T}$, prove that
- $P'(t) = \frac{1}{T} P(t)$
 - $\frac{d}{dt} \ln P(t) = \frac{1}{T}$
9. Plot a graph of $N(t)$ vs t over the interval $0 \leq t \leq 18$ hours if $N(t) = N_0 e^{-\lambda t}$, where $N_0 = 10^{20}$ and $\lambda = 6.4 \times 10^{-5} \text{ s}^{-1}$.
- on linear paper
 - on log-linear paper.
10. (a) Make a table of values of reactor power $P(t)$ and linear rate, $P'(t)$ with 20-second increments in t over the interval $0 \leq t \leq 5$ minutes. Assume $P_0 = 100 \text{ W}$ and $\frac{\Delta k}{L} = 0.05$. Express P and P' in units of % full power, assuming full power equals 100 MW.
- (b) Show consecutive positions of indicating needles on the following meters, at 20-second intervals.





- (c) Describe the needle's motion across each of the above scales, and relate descriptions to the mathematical expressions for linear rate and rate log power.
- (d) Which meter is more suitable for monitoring power at low power levels? At high power levels?
- (e) Which of the following signals is more appropriate for reactor power control
 - (i) at low power levels?
 - (ii) at high power levels?

a signal whose output is proportional to reactor power P , or one whose output is proportional to the logarithm of reactor power, $\log P$?

- 11. Explain the advantage of a rate log signal for reactor protection.
- 12. Show that $\frac{d}{dt} (\log P(t)) = \frac{\Delta k}{L} \log e$, where $\log P$ is the common logarithm of P .

L.C. Haacke